## **Abstract**

The Cambrian age khewra formation sandstone is petrographically analyzed in this research.

Detail mineralogy, texture and diagenesis phenomena were studied. The quartz remains the dominant framework mineral in all sections while feldspar were the second dominant framework grain. Khewra Sandstone is scarce of lithics fragment revealed in this petrographic analysis. The percentage of matrixremains higher throughout in all sections and appeared to be directly related to higher percentages of feldspar as the alteration of feldspar to clay was observed in most of the thin section it accounts for higher percentage of matrix. The color, alteration and replacement of minerals and different orientation of grains narrates that the Khewra Sandstone subjected to all three diagenetic stages. The Khewra Formation lies in sub-arkosic class.

## Acknowledgement

It is a moment of great satisfaction and we are deeply thankful to Almighty ALLAH, Who enabled us to successfully complete the laborious job of this thesis. We are extremely thankful to our Holy Prophet Hazrat Muhammad (P.B.UH) who is a symbol for humans in every field of life.

We wish to express our sincere and deep sense of gratitude to our thesis supervisor Mr. Hummad Ghani Lecturer Bahria University Islamabad, who was very kind and helpful to us at every moment. Through his guidance and kind advice we have completed this task successfully.

We are thankful to Dr. Muhammad Zafar, HOD Department of Earth and Environmental Sciences, Bahria University Islamabad for providing us lab facilities.

We are deeply appreciate the kind support of Mr. Ishtiaq Noor, Senior Geologist HDIP, Islamabad, Mr. Saqib, Lecturer Bahria University Islamabad, Mr. Rashid for sharing their precious time and for giving us useful advices to complete this work successfully. Further sincere support of Mr. Rasheed Maseeh, Rock cutting lab assistant, during the preparation of thin sections is duly acknowledge.

Finally, we are thankful to all our friends specially, Mr. Umer Farooq, Mr. Shuja Khan, Mr. Qaiser Hussain, Mr. Wajid Ali, Mr. Mehmood Ahmad Khan, Mr. Raheem Ullah, Mr. Ijaz Khan, Mr. Ahmad, Mr. Noortaj Khan for their kind and volunteer support during field and labwork.

Sincere acknowledgement to our brothers, sisters, cousins, for their moral support and motivating attitude to complete a laborious task for this thesis.

We are grateful to our parents who prayed a lot for us and provided us moral and financial support.

i

## **Table of Contents**

Acknowledgmenti
Abstractii
Chapter 11
Introduction1
1.1. Location and Accessibility
1.2. Climate and Geography of the Area
1.3. Previous Work
1.4. Aims and Objectives
1.5. Methodology
Chapter 25
Tectonics of the area5
2.1. Regional structural setting6
2.2. Salt Range8
2.2.1. Structure of Salt Range9
2.2.1.1. Compressional Deformation
2.2.1.2. Transform Deformation
2.2.1.3. Extensional Deformation
iii
Chapter 311
Stratigraphy11
3.1. Salt Range Formation (Precambrian)
3.2. Jhelum Group
3.2.1. Khewra formation
3.2.2. Kussak formation
3.2.3. Jutana formation
3.2.4. Baghanwala Formation10
3.2.5. Tobra Formation1
Chapter 424

Petrography	24
4.1. Mineralogy	24
4.1.1. Modal Mineralogy	24
4.1.2. Framework components	24
4.1.2.1. Quartz	25
4.1.2.2. Feldspar	25
4.1.2.3. Lithic grains	25
4.1.2.4. Accessory minerals	26
4.1.2.5. Matrix	26
iv	
4.1.2.6. Cement	26
4.2. Texture	32
4.2.1.Fabric	32
4.2.2. Roundness and Sphericity	33
4.2.3. Sorting	33
4.2.4. Maturity	34
4.3. Classification	36
4.3.1. Sandstone Classification	36
4.3.2. Classification of Khewra Sandstone	37
4.4. Summary	39
Chapter 5	40
Diagenesis	40
5.1. Stages of diagenesis	40
5.2. Diagenesis of Khewra Formation	42
Conclusion	46
References	47

List of	Figures
errains of	Pakistan
Dolzieton	charring

List of Figures
Figure 2.1. Tectono-stratigraphic terrains of Pakistan delineated by major fault boundaries6
Figure 2.2. Tectonic Map of north Pakistan, showing major structural features9
Figure 3.1. Field photograph showing slump bedded sandstone with alternating shale21
Figure 3.2. Field photograph showing interbedded sandstone and shale
Figure 3.3. Field photograph showing planar cross-bedding
Figure 3.4. Field photograph showing dark mineral bands in sandstone
Figure 3.5. Field photograph showing cross-bedding in sandstone
Figure 3.6. Field photograph showing parting lineation in sandstone
Figure 3.7. Contact between Khewra Formation and Khussak Formation. Light grey colour
showing Khussak formation
Figure 4.1. A&B(XPL), C(PPL) showing unstrained quartz
Figure 4.3. A (XPL) (40X) arrow shows zircon inclusion in quartz,
B (XPL) (40X) arrow shows polysynthetic twining in albite,
C (XPL) (20X) arrow shows microcline exhibiting cross-hatch twining,
D (XPL) (20X) showing alteration of feldspar to clay29
Figure 4.4. A (PPL) (40X) arrow shows lithic fragment,
B (XPL) (20X) arrow shows metamorphic lithic30
Figure 4.5. A&B (XPL,PPL) (4X) preferred orientation of mica, observed in thin section,
C&D (XPL) (20X) arrow shows muscovite flake among framework grians30
vi
Figure 4.6. A (XPL) (10X) arrow shows muscovite flake in thin section, B (XPL) (10X) arrows
shows biotite
C&D (XPL,PPL) (20X) arrow shows brown tourmaline as accessory mineral31
Figure 4.7. A&B (XPL,PPL) (4X) arrows shows opaque minerals in thin sections,
C (XPL) (4X) arrows shows matrix among the frame work grain32
Figure 4.8. A&B (XPL,PPL) (20X) arrow shows calcite cement in thin sections32
Figure 4.9. A&B (XPL,PPL) (20X) arrow shows concavo-convex contacts in thin sections
C (XPL) (20X) arrow shows long and sutured contact
Figure 4.10. A&B (XPL,PPL) (4X) preferred orientation in muscovite

Figure 4.11. Sandstone classification scheme for Wackes after Pettijohn, (1987). Q (quartz),
F (feldspars), RF (rock fragments)38
Figure 4.12. classification scheme for Sandstone after Pettijohn, (1987). Q (quartz),
F (feldspars), RF (rock fragments)39
Figure 5.1. A (XPL) (20X) arrow shows bleached or buff sandstone,
B (XPL) (20X) arrow shows calcite cement,
C (XPL) (20X) arrow shows replacement of calcite
Figure 5.2. A (XPL) (20X) arrow shows alteration of feldspar (Alt Fed) to clay minerals.
B(XPL) (20X) arrows shows fractured quartz45
vii
Figure 5.3. A&B (XPL,PPL) (20X) arrow shows concavo-convex contact,
C (XPL) (20X) arrow shows long, concavo-convex and sutured contact46
Figure 5.4. A&B (XPL, PPL) (4X) crystallized mica along grain boundary
List of tables

Table 1. Detailed mineralogical composition of the studied samples......52

Table 2. Normalized values of Quartz (Q(N), Feldspar (F(N), and Lithics(L(N)
Tables 3. Types of contacts observed in thin sections
List of Chart
Chart 1. General Stratigraphic chart of the area
Chart 2. Textural properties of studied thin-sections55
List of Figures Figure 1.1. Accessibility Map of Khewra Gorge (study area)
Figure 2.1. Tectonic Map of north Pakistan, showing major structural features (After Kazmi and
Rana,
1982)
Pakistan, box showing the study area Khewra gorge (Gee, 1989)
8
Figure 3.1. Field photograph showing slump bedded sandstone with alternating shale20
Figure 3.2. Field photograph showing interbedded sandstone and shale.20Figure 3.3. Field photograph showing planar cross-bedding.21Figure 3.4. Field photograph showing dark mineral bands in sandstone.21Figure 3.5. Field photograph showing cross-bedding in sandstone.22Figure 3.6. Field photograph showing parting lineation in sandstone.22Figure 3.7. Base of Khewra Formation consisting of maroonish shale and
sandstone23
Figure 3.8. Contact between Khewra Formation and Khussak Formation shown with black
arrow
Figure 4.1. A&B(XPL), C(PPL) showing unstrained quartz
B (XPL) (40X) arrow shows polysynthetic twining in albite,
C (XPL) (20X) arrow shows microcline exhibiting cross-hatch twining,
D (XPL) (20X) showing alteration of feldspar to clay28
vi
Figure 4.4. A (PPL) (40X) arrow shows lithic fragment,
B (XPL) (20X) arrow shows metamorphic lithic29
Figure 4.5. A&B (XPL,PPL) (4X) preferred orientation of mica, observed in thin section,

C&D (XPL) (20X) arrow shows muscovite flake among framework grians29
Figure 4.6. A (XPL) (10X) arrow shows muscovite flake in thin section, B (XPL) (10X) arrows
shows biotite
C&D (XPL,PPL) (20X) arrow shows brown tourmaline as accessory mineral30
Figure 4.7. A&B (XPL,PPL) (4X) arrows shows opaque minerals in thin sections,
C (XPL) (4X) arrows shows matrix among the frame work grain31
Figure 4.8. A&B (XPL,PPL) (20X) arrow shows calcite cement in thin sections31
Figure 4.9. A&B (XPL,PPL) (20X) arrow shows concavo-convex contacts in thin sections
C (XPL) (20X) arrow shows long and sutured contact
Figure 4.10. A&B (XPL,PPL) (4X) preferred orientation in muscovite35
Figure 4.11. Sandstone classification scheme for Wackes after Pettijohn, (1987). Q (quartz),
F (feldspars), RF (rock fragments)37
Figure 4.12. classification scheme for Sandstone after Pettijohn, (1987). Q (quartz),
F (feldspars), RF (rock fragments)38
vii
Figure 5.1. A (XPL) (20X) arrow shows bleached or buff sandstone,
B (XPL) (20X) arrow shows calcite cement,
C (XPL) (20X) arrow shows replacement of calcite
Figure 5.2. A (XPL) (20X) arrow shows alteration of feldspar (Alt Fed) to clay minerals.
B(XPL) (20X) arrows shows fractured quartz44
Figure 5.3. A&B (XPL,PPL) (20X) arrow shows concavo-convex contact, C (XPL) (20X) arrow
shows long, concavo-convex and sutured contact
Figure 5.4. A&B (XPL, PPL) (4X) crystallized mica along grain boundary45