

Factors Associated With Timely Initiation Of Breastfeeding Among Mothers In JPMC

Beena Barkat Ali, Shazia Naseeb, Razia Korejo

Objective: Study the frequency of common factors Associated with timely initiation of breastfeeding.

Study design: Descriptive Cross sectional.

Duration and place of study: this study was conducted at, Jinnah postgraduate Medical center Karachi unit-I from 19 May to 18 November 2015.

Material & Methods: A total of 108 mother infant pairs were selected by Non probability consecutive sampling technique, meeting our inclusion criteria. Informed consent was taken after explaining the pros and cons, purpose and procedure of the study. The common factors associated with timely initiation of breastfeeding like age of the mother, gestational age, parity, educational status, working status of the mother, mode of delivery, birth weight of the baby and gender of the baby were evaluated through face to face interview of the mothers .

Results: In our study mean age of mothers found to be 26.1 years old. Mean birth interval found to be 3.1years. Mean birth weight of the babies was 2.86 kg. Mean gestational age of the Patients was 38.1weeks. Most of the babies were female 63(58%) other were male 45(42%).Out of 108 patients, 40(37%) received Primary education, 27 (25%) were graduated, 18(17%) were Illiterate, 13(12%) secondary and 10(9%) were Intermediate. Thirty eight (35.2%) belonged to middle class, 56(51.9%) to upper middle class and only 14(13%) to higher class. Fifty eight (53.7%) women were multiparae and 76(70.3%) were employed; And Mostly 66 (61%) delivered through Cesarean -Section.

Conclusion: According to this study the main conclusion of Common factors associated to timely initiation of breastfeeding were age of mothers(74.07%) more than 20years , parity 2 or more(61.1%), educational level secondary and above (50.9%), house wives (32.9%), male sex of infant (41.6%) and Vaginal mode of delivery (38.8%)

Keywords: Breastfeeding, Breastfeeding initiation, primigravidae, practices

Introduction:

Many women want to breastfeed but are unable to do so. Lack of confidence in their ability to breastfeed, problems with proper positioning of neonate, myths of inadequate milk supply, breast pain and lack of support from health professionals in early post discharge period are some reasons why breastfeeding is not initiated and if initiated is not continued for the recommended duration².

Antenatal counseling and postnatal lactation support, improve rates of exclusive breastfeeding. There has been a substantial improvement over the past two decades in the proportion of mothers receiving antenatal care from a skilled health provider, increasing from 26% in 1990-91 to 61 percent in 2006-07 and 73% in 2012-13. Antenatal visits provide an opportunity to educate women regarding the benefits of

breastfeeding and can help improve rates of initiation of exclusive breastfeeding¹⁻⁴.

Baby Friendly Hospital Initiative was launched in 1991 by UNICEF⁴ and WHO⁵ to ensure that all maternity services support breastfeeding. Ten steps to successful breastfeeding need to be implemented if a health facility wants to be accredited as Baby Friendly. Step 3 of these steps is 'inform all pregnant women about benefits and management of breastfeeding'. Implementation of baby friendly hospital initiative in Sindh, Pakistan improved breastfeeding practices in some of the centers to 98.97%⁵.

Breastfeeding practices in Pakistan are far from ideal. Ninety-four percent of children were reported to have been breastfed at some time. 38% percent of children less than age 6 months are exclusively breastfed. The median duration of exclusive breastfeeding is less than one 1month. Complementary foods are not introduced in a timely fashion for all children. Only 57% of breastfed children age 6-9 months received complementary foods. Overall, only 15% of children ages 6-23 months are fed appropriately based on recommended infant and young child feeding (IYCF) practices².

Breastfeeding offers many advantages to the newborn and the new mother. Although emphasis is put on breastfeeding rates, even then, breastfeeding often fall short of the required frequency of breastfeeding.

The WHO⁶ and American Academy of Pediatrics⁷ both recommend exclusive breastfeeding for six months and

Beena Barkat Ali,
Postgraduate Student,
Jinnah Post Graduate Medical Center, Karachi

Shazia Naseeb,
Associate Professor, Dept. of Gynaecology & Obstetrics
Jinnah Post Graduate Medical Center, Karachi
Email:snkhan1975@yahoo.com

Razia Korejo,
Professor, Dept. of Gynaecology & Obstetrics
Bahria University Medical & Dental College, Karachi

Received: 20-02-2018
Revised: 30-04-2018
Accepted: 15-05-2018

complementary feeding with breastfeeding for at least 12 or 24 months. In our study 78% of women who were counselled knew that exclusive breastfeeding needs to be practiced for first six months and only 35% knew that babies less than 6 months do not need extra water. Ahmad et al report 68% of women breastfeed exclusively after breastfeeding and Dhandapany et al report that more women practice exclusive breastfeeding after counselling^{8,9}

The Pakistan Demographic and Health Survey² also reported introduction of complementary feeding in 10% of infants less than 6 months and 19% of infants aged 4-5 months. Bottle-feeding is reported as a norm even though not supported by health professionals. More than 1 in 5 babies under two months of age are bottlefed³. Similar findings could not be gathered from our study, as the patients were not followed up in the postnatal period. Our study was limited in this context. Early initiation of breast feeding, especially within one hour of birth, refers to the best practice recommended by WHO¹⁰. Early initiation of breast feeding will directly support progress toward achieving MDG through reducing neonatal mortality¹¹.

In Pakistan breastfeeding is nearly universal although early initiation is not common. A survey done in 1990-91 revealed that only 8.5% of neonates were breastfed within the first hour and only 25.8% were breastfed on the day of delivery. The estimates rose to 27.2% and 65.5% in 2006-2007, shows considerable improvement¹².

The rationale of the study is that on extensive literature search study on factors associated with timely initiation of breast feeding has not been done locally and there is dearth of literature internationally as well. Therefore this study is designed to estimate the magnitude of these factors and to generate local data. Secondly some policy could be devised to highlight this issue thereby timely initiation of breast feeding could be initiated that will prevent maternal and neonatal morbidity and mortality.

Material and Method

Women meeting the inclusion criteria admitted in the Department of Obstetrics and Gynecology, JPMC, Karachi, were enrolled in the study. It was a cross sectional study. Mothers were selected through non probability technique. Informed consent was taken after explaining the pros and cons, purpose and procedure of the study. The factors associated with timely initiation of breastfeeding like age of the mother, gestational age, parity, educational status, and occupation of the mother, mode of delivery, birth weight of the baby and gender of the baby were evaluated through face to face interview of the mothers and confirmed by hospital records and documented on structured Performa .

SPSS version 20 was used for data entry and analysis. Mean±SD was calculated for age of the women and gestational age at delivery, birth weight of the baby and birth interval. Frequency and percentages were calculated

for gender of the baby, parity, educational level, mode of delivery, occupation of women, family monthly income and family structure (nuclear or extended). Effect modifiers were controlled through stratification of family monthly income, family structure (nuclear or extended) and birth interval to determine the effect of these on outcomes. Chi square test were applied and p value =0.05 was taken as significant.

Results:

Mean age of the patients was 26.1 years ranging from (18-35) years. Mean birth interval of the baby was 3.1years. Mean birth weight of the baby was 2.8 Kg. Mean gestational age of the Patients was 38.1 weeks. The gender distribution of the baby most of the babies were female 63(58%) other were male 45(42%). Out of 108 mothers 18(17%) were Illiterate, 40(37%) were Primary educated, 13(12%), were secondary, 10(9%) were Intermediate and 27 (25%) were graduated. The distribution of monthly income of mother women 38(35.2%) belong to middle class (monthly income <25,000 RS.), 56(51.9%) to upper middle class (monthly income 25,000-50,000 RS.) and only 14(13%) belong to higher class (monthly income >50,000 RS.). Fifty-Eight (53.7%) women had multiparty. Most of the women were employed 76(70.3%). Forty-two patients (38.9%) women belonged to age 26-30 years with next majority between 21-25 years i.e. 36 (33.3%) and only 5(4.7%) patients were more than 30 years of age. The gestational ages at delivery ranged from 34 to 42 weeks but most of them belong to gestational age more than 37 weeks i-e 80(74%). Most of the women 66(61%) who have delivered baby through C-Section. Effect modifiers were controlled through stratification of family monthly income, family structure (nuclear or extended) and birth interval to determine the effect of these on outcomes. Chi square test was applied and p value =0.05 was taken as significant as shown on (Table-2-3). There was significant association is seen family income and gender together affecting breast feeding(table-2), significant association is seen family income and Parity, family income and gest.age affecting breast feeding(Table-3) and significant association is also seen family structure and education, family structure and mode of delivery affecting breast feeding(Table-4).and birth interval and parity, birth interval and gest. Age are also showing significant association.

Discussion:

Breastfeeding is advocated as an important child survival strategy by the World Health Organization especially in countries with poor socioeconomic background¹⁻². Breast-feeding is fundamental to the health and development of children and important for the health of their mother as well.

In Pakistan breastfeeding is nearly universal although early initiation is not common. A survey done in 1990-91 revealed that only 8.5% of neonates were breastfed within the first hour and only 25.8% were breastfed on the day of delivery. The estimates rose to 27.2% and 65.5% in 2006-2007, shows

CHARACTERISTICS OF MOTHERS	FREQUENCY	PERCENTAGES (%)
Age		
<20 Years	25	23.1%
21-25 Years	36	33.3%
26-30 Years	42	38.9%
>30 Years	5	4.7%
Parity		
Primiparae	42	38.9%
Multiparae	58	53.7%
Grand multiparae	8	0.4%
Educational Status		
Illiterate	18	17%
primary	40	37%
secondary	13	13%
Intermediate	10	9%
Graduate	27	25%
Gender of Baby		
Male	45	42%
Female	63	58%
Birth Weight of baby		
<3kg	35	32%
>3kg	73	68%
Mode of Delivery		
Vaginal delivery	42	39%
Cesarean- sections	66	61%
Profession of Mothers		
House wife	32	29.7%
Working	76	70.3%

Table 1. Demographic and Obstetric Characteristics of Mothers n=108

	Age more than 20 years	Age less than 20 years	Total	P-value	Baby Gender Male	Female	Total	P-value
Family Income Rs < 25,000 Rs 25-50,000 Rs >50,000	20(18.5%) 47(43.5%) 13(12%)	18(16.7%) 09(8.3%) 1(0.9%)	38(35.2%) 56(51.9%) 14(13%)	0.001*	30(27.8%) 10(9.3%) 05(4.6%)	8(7.4%) 46(42.6%) 09(8.3%)	38(35.2%) 56(51.9%) 14(13%)	0.000*
Family Structure Nuclear Extended	35(33%) 45(42.5%)	03(2.8%) 23(21.7%)	38(35.8%) 68(64.2%)	0.004*	16(14.8%) 29(26.9%)	29(26.9%) 34(31.5%)	45(41.7%) 63(58.3%)	0.276
Birth Interval 1-2 3-4 >5	45(41.7%) 35(32.4%) 0(0%)	16(14.8%) 10(9.3%) 2(1.9%)	38(35.8%) 68(64.2%) 68(64.2%)	0.049	25(23.1%) 13(12%) 2(1.9%)	36(33.3%) 32(29.6%) 0(0%)	61(56.5%) 45(41.7%) 2(1.9%)	0.079

Table 2. Association of Family Income and Gender affecting breast feeding

	Parity 2 &>2	Parity <2			Gest.Age Full term baby 37-42	<37 Weeks		
Family Income Rs < 25,000 Rs 25-50,000 Rs >50,000	15(41.7%) 39(18.5%) 12(0.9%)	23(14.8%) 17(23.1%) 2(0.9%)	38(35.2%) 56(51.9%) 14(13%)	0.002*	28(25.9%) 46(42.6%) 06(5.6%)	10(9.3%) 10(9.3%) 08(7.4%)	38(35.2%) 56(51.9%) 14(13%)	0.011*
Family Structure Nuclear Extended	25(23.1%) 41(38%)	15(13.9%) 27(25%)	45(37%) 68(63%)	0.820	30(27.8%) 50(46.3%)	15(13.9%) 13(12%)	45(41.7%) 63(58.3%)	0.276
Birth Interval 1-2 3-4 >5	45(41.7%) 20(18.5%) 1(0.9%)	16(14.8%) 25(23.1%) 1(0.9%)	61(56.5%) 45(41.7%) 2(1.9%)	0.009*	51(47.2%) 29(26.9%) 0(0%)	36(9.3%) 32(14.8%) 2(1.9%)	61(56.5%) 45(41.7%) 2(1.9%)	0.005*

Table 3. Association Family income with Parity of the Mothers and Gestational age

	Education Level: secondary & above	Primary or less			Spontaneous vaginal delivery	C section		
Family Income Rs < 25,000 Rs 25,50,000 Rs >50,000	23(21.3%) 30(27.8%) 05(4.6%)	15(13.9%) 26(24.1%) 9(8.3%)	38(35.2%) 56(51.9%) 14(13%)	0.282	15(13.9%) 24(22.4%) 03(2.8%)	23(21.3%) 33(30.6%) 10(9.3%)	38(35.2%) 56(51.9%) 14(13%)	0.445
Family Structure Nuclear Extended	30(27.8%) 20(18.5%)	15(13.9%) 43(39.8%)	45(41.7%) 63(58.3%)	0.000*	29(26.9%) 13(12%)	16(14.8%) 50(46.3%)	45(41.7%) 63(58.3%)	0.000*
Birth Interval 1-2 (years) 3-4 >5	40(37%) 10(9.3%) 0(0%)	21(19.4%) 35(32.4%) 2(1.9%)	61(56.5%) 45(41.7%) 2(1.9%)	0.000*	24(22.2%) 17(15.7%) 1(0.9%)	10(34.3%) 16(25.9%) 1(0.9%)	61(56.5%) 45(41.7%) 2(1.9%)	0.936

Table 4. Association of Family structure, birth interval and mode of delivery

considerable improvement¹².

Promotion of early initiation of breast feeding is necessary for the prevention of avoidable deaths of children as different studies have proved that death rate is high when breast feeding started late¹³. It is shown from different studies that factors like antenatal visits, knowledge about breast feeding, number of live births, age of mother, mode of delivery, her education, occupation and economical status all have strong association with early initiation of breast feeding.

Breastfeeding is multi-factorial in nature and different factors will be at play depending on individual circumstances. We found similar results when we compared our study with a study which was done at Saudi Arabia in that study. The independent predictors of timely initiation were mothers who refrain from prelacteal feeding (16%) and mother's residence rural area (19.2%), absence of breast problems (12.7%) and parity 2 Or more 27%.10 other factors were age >20 years (11.34%), educational level secondary and above (22%), house wife (11.8%), male sex of infant (13.4%). Full term (12.6%), mode of delivery; spontaneous vaginal delivery (13%)¹⁴.

Sharma A et al¹⁵ also found significant association similar to our study between early initiation of breast feeding with education, occupation and economic status of the mother but reported no significant association with maternal age,

type of family, family size and live birth. Setegn et al¹⁶ also reported same as formal educated women were 1.4 times as likely to initiate breast feeding with first hour of delivery.

In many developing countries particularly in Asia, the converse is true and extended lactation is more frequent in poor uneducated women in rural areas, and in urban areas higher level of education are negatively associated with breast feeding initiation and duration¹⁷⁻¹⁸.

Other studies in Jeddah, Saudi Arabia by Fida and Al-Aama (2003) and Shawky and Abalkhail (2003)¹⁹⁻²² included 21% of university graduated women (Fida & Al-Aama, 2003), 39.5% illiterate women and 88% housewives (Shawky & Abalkhail, 2003)²²⁻²⁵. A more recent study, which also was done in a private hospital in Jeddah, Saudi Arabia (Mosalli et al, 2012)²³ reported fewer multiparous (54%), graduated (24%), and caesarean delivery (31%) women compared with current study. However, Mosalli et al. (2012) reported similar percentages of housewives (78%) and similar age group distribution.

Many studies have indicated that a woman's obstetric experience may influence her breast-feeding behaviors. Our study shows positive association with vaginal delivery while some researchers²⁴ have found no association between mode of delivery and breastfeeding, but others²⁵ have reported a negative association between cesarean delivery and breast-

Factors Associated With Timely Initiation Of Breastfeeding Among Mothers In JPMC

feeding initiation but not duration once breast feeding has commenced.

Conclusion:

Hence we conclude that age of mother (being more than 20 years), having two or more children, having at least secondary level of education, having male last child, having last birth by spontaneous vaginal delivery and being a house wife were factors responsible for timely initiation of breast feeding in our study.

Recommendation:

In the light of our findings we recommend that breastfeeding education should be in conjunction with obstetrics. Hence two clinics should be merged into one at Jinnah Postgraduate Medical Center Karachi.

REFERENCES:

1. Millenium Development Goals. Available from:http://www.who.int/topics/millennium_development_goals/en/ Accessed on May 29 201.
2. National Institute of Population Studies. Pakistan Demographic and Health Survey 2012-2013[Internet]. Islamabad: National Institute of Population Studies; 2013. Available from: <http://www.nips.org.pk>. Accessed on November 25,2014.
3. Suresh S, Sharma KK, Saksena M, Thukral A,Agarwal R, Vatsa M. Predictors of breastfeeding problems in the first postnatal week and its effect on exclusive breastfeeding rate at six months: experience in a tertiary care centre in Northern India. *Indian J Public Health* 2014; 58: 270-3.
4. The Baby Friendly Hospital Initiative [Internet].UNICEF. Available from: <http://www.unicef.org/programme/breast-feeding/baby.htm>. Accessed on December 5, 2014.
5. Khan M, Akram DS. Effects of baby-friendly hospital initiative on breast-feeding practices in sindh. *J Pak Med Assoc* 2013; 63(6): 756-9
6. World Health Organization. The optimal duration of exclusive breastfeeding: report of an expert consultation Geneva: WHO, 2001. [http:// www.who.int/nutrition/publications/optimal_duration_of_exc_bfeeding_report_eng.pdf](http://www.who.int/nutrition/publications/optimal_duration_of_exc_bfeeding_report_eng.pdf). Accessed on 29 May 2015
7. Gartner LM, Morton J, Lawrence RA, Naylor AJ,O'Hare D, Schanler RJ, et al. American Academy of Pediatrics Section on Breastfeeding. Breastfeeding and the use of human milk. *Pediatrics* 2005; 115(2): 496-506
8. Ahmad MO, Sughra U, Kalsoom U, Imran M, Hadi U. Effect of antenatal counselling on exclusive breastfeeding. *J Ayub Med Coll Abbottabad* 2012; 24(2): 116-9.
9. Dhandapany G, Bethou A, Arunagirinathan A, Ananthkrishnan S. Antenatal counseling on breastfeeding -- is it adequate? A descriptive study from Pondicherry, India. *Int Breastfeed J* 2008; 3(1): 5.
10. WHO Breastfeeding-early initiation :World Health Organization;2012 <http://www.who.int/Elena/title/earlybreast-feeding/en/access> 5 April 2013
11. Khan J,VesellL,BahlR, MartinesJc.,Timing of breastfeeding during first month of life: Effect on neonatal mortality and morbidity-a systemic review and meta analysis:*mat and child nutr.* 2015; 19(3): 468-79.
12. National Institute of Population Studies. Pakistan Demographic and Health Survey 2006-2007 [Internet]. Islamabad: National Institute of Population Studies; 2013.Available from: <http://www.healthkp.gov.pk/downloads/PDHS.pdf> Accessed on 29 may 2015.
13. Santos et al .Avoidable deaths in first four years of life among children in 2004 pelotos(brazil birth cohort study. *ca demos de saude public.*2011; 27: 185-97
14. El-Gilany AH, Sarraf B, Al-Wehady A. Factors associated with timely initiation of breastfeeding in Al-Hassa province, Saudi Arabia. *East Mediterr Health J.* 2012; 18(3): 250-4.
15. Sharma A et al.factors associated with early ignition of breast feeding among mothers of tribal areas of Madhya paradesh:a community based,India:a community based cross sectional study; *int J community med.public health.* 2017; 3(1): 194-9.
16. Setengn T,Gerbaba M, BelachewT.Determinants of timely initiation of breast feeding among mothers in Goba woreda,south east Ethiopia: A cross sectional study.*BMC public health.* 2011; 11(1): 217
17. Penticen.Extended breast-feeding and growth in rural china.*Nute.Rev.* 114; 52: 144-48.
18. Rogers I ,Emmett P,GoldingJ. The incidence and duration of breast feeding.*Early Hum Devel.*1997; 49(Supplement): 143-155.
19. Truman, Edwin M. Sovereign wealth funds: the need for greater transparency and accountability. Peterson Institute for International Economics, 2008.
20. Al-Mazrou, Y. Y., K. M. S. Aziz, and M. Khalil. "Breastfeeding and weaning practices in Saudi Arabia." *Journal of tropical pediatrics.* 1994; 40(5) 267-71.
21. Fida, Nadia M., and Jumama Y. Al-Aama. "Pattern of infant feeding at a University Hospital in Western Saudi Arabia." *Saudi medical journal.* 2003; 24(7); 725-9.
22. Shawky, Sherine, and Bahaa A. Abalkhail. "Maternal factors associated with the duration of breast feeding in Jeddah, Saudi Arabia." *Paediatric and perinatal epidemiology.* 2003; 17(1); 91-6.
23. Mosalli, Rafat, et al. "Perceived barriers to the implementation of a baby friendly initiative in Jeddah, Saudi Arabia." *Saudi medical journal.* 2012; 33(8); 895-900.
24. Scott J A,Landers MCG,Hughes RM,Binns CW.factors associated with breast feeding at Discharge and duration of breast feeding *JPediatar child Health.* 2001; 37(3): 254-261.
25. Perez-EscamillaR, Maulen-RadovanI, Dewey KG. The association between caesarean delivery and breast feeding outcome among maxicon women. *Am J public Health.* 1996: 86(6); 832-6.

